Problems with the "Fundamental Theorem of Algebra".

When students learn about the Fundamental Theorem of Algebra they clearly understand the basic idea but they have many "***But what if?***" questions which make them feel that the theorem only works in special cases.
This article is written for students at this level and for teachers who present them with the concept.
I strongly believe that articles of this type must be written in "*plain language*" at the level of the students it is aimed at.
Basically, the theorem states that equations like:

1. $x^2 - 3x + 2 = 0$ will always have 2 solutions
2. $x^3 - 7x^2 + 14x - 8 = 0$ will always have 3 solutions
3. $x^4 - 5x^2 + 4 = 0$ will always have 4 solutions
4. $x^7 + ax^6 + bx^5 \ldots + c = 0$ will always have 7 solutions
5. $ax^n + bx^{n-1} + cx^{n-2} + \ldots + p = 0$ will always have n solutions

The above equations are called polynomial equations (powers of x) and simply stated, the theorem states that if x^n is the highest power, the equation will have n solutions (where n is a positive integer).

I believe the theorem must be related to graphs to get a clear understanding.

I will go through three straightforward cases where the theorem applies nicely without any ambiguity.

1. Consider the graph of $y = x^2 - 3x + 2$ which is a simple parabola.
If we say "***When is*** $x^2 - 3x + 2 = 0$?" we are saying "***When is*** $y = 0$***?***" which clearly means "***When does the graph cross the*** x ***axis***?"



This is the graph of $y = x^2 - 3x + 2$
which clearly crosses the x axis at
$x = 1$ and at $x = 2$

which means that when $y = 0$
then $x = 1$ and $x = 2$

which also means that when $x^2 - 3x + 2 = 0$
then $x = 1$ and $x = 2$

This has 2 solutions because the highest power is 2.

2. Now consider the graph of $y = x^3 - 7x^2 + 14x - 8$ which is called a cubic curve.

If we say "***When is*** $x^3 - 7x^2 + 14x - 8 = 0$?" we are saying "***When is*** $y = 0$***?***" which clearly means "***When does the graph cross the x axis***?"



This is the graph of $y = x^3 - 7x^2 + 14x - 8$ which clearly crosses the x axis at $x = 1$, at $x = 2$ and at $x = 4$

which means that when $y = 0$ then $x = 1$, $x = 2$ and $x = 4$

which also means that when $x^3 - 7x^2 + 14x - 8 = 0$ then $x = 1$, $x = 2$ and $x = 4$

This has 3 solutions because the highest power is 3.

3. Now consider the graph of $y = x^4 - 5x^2 + 4$ which is called a quartic curve.

As stated before, if we say "***When is*** $x^4 - 5x^2 + 4 = 0$?" we are saying "***When is*** $y = 0$***"*** which clearly means "***When does the graph cross the x axis***?"



This is the graph of $y = x^4 - 5x^2 + 4$ which clearly crosses the x axis at $x = -2$, $x = -1$, $x = 1$ and $x = 2$

which means that when $y = 0$ then $x = -2$, $x = -1$, $x = 1$ and $x = 2$

which also means that when $x^4 - 5x^2 + 4 = 0$ then $x = -2$, $x = -1$, $x = 1$ and $x = 2$

This has 4 solutions because the highest power is 4.

At this point of course we expect such questions as ***"But what if the graph does not cross the x axis? The theorem does not seem to work!"***

The usual "answer" to this type of question is that ***"The solutions are imaginary numbers but the equation still has the right number of solutions"!***

Another common question is when the graph just **touches** the x axis and students are merely told ***"This is called a double solution!"***

These sort of inadequate, unconvincing answers make students lose any respect for the theorem but there is a very nice way to show that the graphs actually do have **physical places** where the ***y*** value is zero!

INTRODUCTION TO "PHANTOM GRAPHS".

Fig 1



This is the basic graph of $y = x^2$ and if we only use **real values** of ***x*** we only obtain **positive** values of ***y***. **Fig 1**

$x = \pm 1$ we get $y = 1$
$x = \pm 2$ we get $y = 4$
$x = \pm 3$ we get $y = 9$

However, we may allow values of ***x*** such as:

$x = \pm i$ we get $y = -1$
$x = \pm 2i$ we get $y = -4$
$x = \pm 3i$ we get $y = -9$

} and these are **real *y*** values!

Fig 2



The **insight**, is to allow a **complex *x* PLANE** but with just a **real *y* AXIS. Fig 2**

This produces a sort of "***phantom***" parabola underneath the basic parabola and at right angles to it.

I have discovered that nearly ALL curves have these extra "***phantom***" parts and more importantly, this has an intriguing connection with the **Fundamental Theorem of Algebra**.

Basically, the **Fundamental Theorem of Algebra** states that polynomial equations of the form: $ax^n + bx^{n-1} + cx^{n-2} + \ldots\ px^2 + qx + r = 0$ will have ***n*** solutions.
(where ***n*** is a positive integer)
This is often interpreted as:
"*The solutions of an equation* ***f(x) = 0*** *are where the graph of* ***y = f(x) crosses the x axis***".
But this only finds the solutions which are REAL numbers! Fig 3



Consider the equation $x^2 - 4x + 3 = 0$
The graph of $y = x^2 - 4x + 3$ is as shown in **Fig 3**

The graph crosses the ***x*** axis at $x = 1$ and $x = 3$ so the solutions are $x = 1$ and 3

In this case, the phantom hanging below had no part to play in this logic.

Fig 4



However, consider the equation $x^2 - 4x + 4 = 0$
The graph of $y = x^2 - 4x + 4$ is as shown in Fig 4.

In this case, the top half of the parabola crosses the ***x*** axis at ***x = 2*** **AND** the bottom half of the parabola (*the phantom*) ALSO crosses the ***x*** axis at ***x = 2. (This is called a "double solution")***

The graph goes through the point (2, 0) <u>TWICE</u>.
The red basic parabola goes through (2, 0) and the purple ***phantom*** parabola also goes through (2, 0)

Fig 5



Of course, the most interesting case is when the basic **top half** of the parabola would not normally cross the ***x AXIS*** at all but its ***phantom* would** cross the ***complex x PLANE***!

Consider the equation $x^2 - 4x + 5 = 0$
The graph of $y = x^2 - 4x + 5$ is as in Fig 5.

The *phantom* crosses the ***x plane*** at
x = 2 + i and x = 2 – i as shown in Fig 5 and these are the complex solutions of the equation.

We can now re-state the **Fundamental Theorem of Algebra** as:
"The solutions, ***whether they are real or complex,*** of an equation ***f(x) = 0,*** are where the graph of ***y = f(x)*** crosses the ***complex x plane***".

Fig 6.



Clearly, we can see that any parabola of the form:
$y = ax^2 + bx + c$ (with its ***phantom***) will cross **any** horizontal plane ***y = c*** (which represents any real ***y*** value) **exactly two times**. **Fig 6**.

CUBIC EQUATIONS

Fig 7.



In my website www.phantomgraphs.weebly.com you will find detailed working to show HOW **CUBIC** functions of the form:
$y = ax^3 + bx^2 + cx + d$
each have **2 phantoms** emanating from their **maximum** and **minimum** points. See **Fig 7**.
(I will show how the equations of the phantoms are calculated later in this article.)

We know that any Cubic equation of the form:
$ax^3 + bx^2 + cx + d = 0$ will have 3 solutions.

Sometimes we have **3 REAL solutions** as for the intersections with the middle **Plane B** in **Fig 7** and sometimes we have **1 real solution and 2 complex solutions** as on **Planes A and C.**
See **Fig 7**.

I will go through these three basic cases for the solutions of cubic equations:

1. Graph showing the three REAL roots of a cubic equation:

$$y = x^3 - x = x(x^2 - 1) = x(x + 1)(x - 1)$$

(Roots are x = 0, x = –1, and x = 1)



In this case the basic RED curve has two turning points from which the two PURPLE phantoms emerge.
Both the upper and lower phantoms do not cross the x plane so they play no part in the solutions of $x^3 - x = 0$
The basic RED curve crosses the x plane THREE times showing the real solutions to $x^3 - x = 0$ ***are x = 0, x = –1, and x = 1***

2. Graph showing one real and two complex roots of the equation:

$$y = x^3 - x^2 + 2 = (x + 1)(x^2 - 2x + 2)$$
$$= (x + 1)\,(x - (1+ i))\,(x - (1 - i))$$
(Roots are $x = -1$, $x = 1 + i$ and $x = 1 - i$)



*In this case the basic **RED** curve has two turning points from which the two **PURPLE** phantoms emerge.*

The upper phantom does not cross the x plane so this does not show any solutions to $y = 0$.

The lower phantom crosses the x plane showing the solutions $x = 1 + i$ and $x = 1 - i$.
Notice that because of the positions of the phantom graphs, they will always produce roots which are complex conjugates (ie $a + ib$ and $a - ib$)

The basic curve crosses the x plane showing the real solution at $x = -1$.

3. Graph showing one real and two complex roots of the equation:

$$y = x^3 - 2x^2 + x - 2 = (x^2 + 1)(x - 2)$$
$$= (x + i)(x - i)(x - 2)$$

(Roots are $x = i$, $x = -i$ and $x = 2$)



*In this case the basic **RED** curve has two turning points from which the two **PURPLE** phantoms emerge.*

The lower phantom does not cross the x plane so this does not show any solutions to $y = 0$.

The upper phantom crosses the x plane showing the solutions at $x = i$ and $x = -i$.

The basic curve crosses the x plane showing the real solution at $x = 2$.

QUARTIC EQUATIONS

Fig 8.



This is a typical **QUARTIC** graph. **Fig 8** showing 1 maximum and 2 minimum points.

This is the same Quartic graph with its 3 *phantoms* emanating from each turning point. **Fig 9**.
The intersection points with the 4 planes A, B, C and D with the graph, are marked.
Plane A shows 2 real solutions on the basic RED curve and 2 imaginary solutions on the PURPLE *phantom.*



Plane B shows 4 real solutions on the basic RED curve.

Plane C shows 2 double real solutions which lie on the basic red curve and the GREEN *phantoms*.

Plane D shows 4 imaginary solutions on the two GREEN *phantoms*. (2 sets of conjugate roots)

(I will show how the equations of the phantoms are calculated later in this article.)

Clearly a **QUARTIC** curve will pass through ANY horizontal plane 4 times.

I will now go through the basic cases for the solutions of quartic equations:

1. Graph showing the 4 REAL roots of a quartic equation:



The basic red graph clearly crosses the ***x axis*** four times.

2. Graph showing a quartic with no real roots but 4 complex roots.



The basic red graph clearly does not cross the ***x axis*** but the two lower ***phantoms*** do cross the ***complex x plane*** 4 times.

3. Graph showing a quartic with 2 sets of double real roots.



The basic red graph is clearly "***sitting***" on the ***x*** axis at 2 points but notice that the two lower phantoms also go through the same points.

The graph does cross (or touch) the ***x*** plane a total of 4 times but we sometimes say, it has 2 "double" solutions.

4. Graph showing a quartic with 2 distinct real roots and a double real root at x = 0.



The basic red graph is clearly "***sitting***" on the underside of the ***x*** axis at ***x*** = 0 and the top phantom is sitting on the top side of the ***x*** axis at ***x*** = 0 also. This is a "double" solution.

The basic red graph also crosses the ***x*** axis at two distinct points on the ***x*** axis.

5. Graph showing a quartic with 2 real roots and 2 imaginary roots.



The basic red graph crosses the ***x*** axis twice which means there are 2 real roots.

The top phantom crosses the ***x*** plane (actually on the imaginary ***x*** axis) which means there are 2 imaginary roots also.

Finding the equations of the *Phantom Graphs*.

This would be a good stage to show the process for finding the equations of the ***phantom graphs*** (which use ***complex x values*** to still produce ***real y values***).

Consider the graph of $y = x^4$
As you can see this is a very U-shaped graph.



Notice that if $x^4 = 16$ then $x = 2$ *and* $x = -2$

But if $x^4 = 16$ then there should be 4 solutions according to the Fundamental Theorem of Algebra but we can only see two solutions here.
$x = 2$ *and* $x = -2$

Also, how can we have 4 solutions to the equation $x^4 = 0$?

I will address these problems now!

We are going to allow some imaginary ***x*** values so we need an imaginary ***x axis*** as shown below.



Example:
The point (2 + 3i, 0)
will be at this position.

Using a 3D coordinate system the point is really (2, 3, 0)

NB
Start from the origin
go 2 units along the real ***x*** axis then
go 3 units along the imaginary ***x*** axis
then don't move up because ***y*** = 0
The point is ON the complex ***x*** plane

Also I will replace the real variable "***x***" by "***x + iz***" so the equation becomes:
$y = (x + iz)^4$
Expanding we get:
$y = (x + iz)^4$
$y = x^4 + 4x^3zi + 6x^2z^2i^2 + 4xz^3i^3 + i^4z^4$
Separating into real and imaginary parts this becomes:
$y = x^4 - 6x^2z^2 + z^4 \quad + \quad i(4x^3z - 4xz^3)$ ———*Equation A*
The important step here is to realise that we only want the complex *x* values which produce REAL *y* values.
This means the **imaginary part of *y* has to be zero** in *Equation A* above!
So $4x^3z - 4xz^3 = 0$
$4xz(x^2 - z^2) = 0$
This means $z = 0$ ***OR*** $x = 0$ ***OR*** $z^2 = x^2$

If $z = 0$ ***then substituting in*** *Equation A*
we get $y = x^4$ ***which is the basic equation in the x, y plane.***

If $x = 0$ ***then substituting in*** *Equation A*
we get $y = z^4$ ***which is a phantom graph the same shape as*** $y = x^4$ ***but in the z, y plane perpendicular to*** $y = x^4$ ***which is in the x, y plane.***

If $z^2 = x^2$ ***which means*** $z = \pm x$ ***then*** $y = x^4 - 6x^4 + x^4 = -4x^4$
Surprisingly, this represents 2 more phantoms the same shape as $y = x^4$ ***but rotated through 45 degrees.***

This is the full graph of $y = x^4$ for <u>real</u> y values (but allowing <u>complex</u> x values).



Notice: I have marked the 4 solutions of $x^4 = 16$ on the top of the graph
x = – 2 ***x = 2i*** ***x = –2i*** ***x = 2***
And if we consider the equation $x^4 = 0$ we see that all 4 branches of the curve do go through $x = 0$.

Recall I said earlier that a more meaningful way to think of the theorem is:
"The solutions, ***whether they are real or complex,*** of an equation ***f(x) = 0,*** are where the graph of ***y = f(x)*** crosses the ***complex x plane***".

For the case of $x^4 = 0$ the graph $y = x^4$ ***does cross the complex x plane 4 times at x = 0.*** It certainly does sound silly but the equation has 4 "solutions" or actually, "intersections" and each solution or intersection is at $x = 0$.

I will also consider the graph of $y = x^3$ and find the equations of its phantoms.



Notice that if $x^3 = 8$ then $x = 2$

But if $x^3 = 8$ then there should be 3 solutions according to the Fundamental Theorem of Algebra but we can only see 1 solution here at $x = 2$

Also, how can we have 3 solutions to the equation $x^3 = 0$?

I will address these problems soon!

Also I will replace the real variable x by $x + iz$ so the equation becomes:
$y = (x + iz)^3$ because this allows for complex x values but only real y values.
Expanding we get:

$$y = (x + iz)^3$$
$$y = x^3 + 3x^2zi + 3x\,z^2i^2 + i^3z^3$$

Separating into real and imaginary parts this becomes:

$$y = x^3 - 3x\,z^2 + i(3x^2z - z^3) \quad \text{——— } \textit{Equation A}$$

The important step here is to realise that we only want the complex x values which produce REAL y values.
This means the **imaginary part of y has to be zero** in *Equation A* above!
So $3x^2z - z^3 = 0$

$$z(3x^2 - z^2) = 0$$

This means $z = 0$ ***OR*** $z^2 = 3x^2$

If $z = 0$ ***then substituting in*** *Equation A*
we get $y = x^3$ ***which is the basic equation in the x, y plane.***

If $z^2 = 3x^2$ ***which means*** $z = \pm x\sqrt{3}$ ***then*** $y = x^3 - 3x3x^2 = -8x^3$
Surprisingly, this represents 2 more phantoms the same shape as $y = x^3$ ***but rotated through 120 degrees to each other.***

This is the full graph of $y = x^3$ for <u>real</u> y values (but allowing <u>complex</u> x values).



Notice: I have marked the 3 solutions of $x^3 = 8$ on the top of the graph

$x = -\frac{1}{2} + i\frac{\sqrt{3}}{2}$ $x = -\frac{1}{2} - i\frac{\sqrt{3}}{2}$ $x = 2$

And if we consider the equation $x^3 = 0$ we see that all 3 branches of the curve do go through $x = 0$.

For the case of $x^3 = 0$ the graph $y = x^3$ ***does cross the complex x plane 3 times at*** $x = 0$.

It is nice to know that these ***phantoms*** seem to appear at the turning points but some curves do not have any turning points!
For instance consider a cubic with roots $x = 1, 1 + i$ and $1 - i$
The equation would be $y = (x - 1)(x - (1 + i))(x - (1 - i))$

$$= (x - 1)(x^2 - 2x + 2)$$
$$= x^3 - 2x^2 + 2x - x^2 + 2x - 2$$
$$= x^3 - 3x^2 + 4x - 2$$

The basic 2D graph looks like this:



It is not at all apparent where the ***phantoms*** could be!

If we allow complex x values we replace x ***by*** $(x + iz)$
The equation is now $y = (x + iz)^3 - 3(x + iz)^2 + 4(x + iz) - 2$
Expanding:

$$y = \left[x^3 - 3xz^2 - 3(x^2 - z^2) + 4x - 2\right] + i\left[(3x^2z - z^3) - 6xz + 4z\right]$$

but we only want ***real y values so*** $3x^2z - z^3 - 6xz + 4z = 0$
factorising***:*** $z(3x^2 - z^2 - 6x + 4) = 0$
this means $z = 0$ ***(which produces the basic 2D graph) ie*** $y = x^3 - 3x^2 + 4x - 2$
or $z^2 = 3x^2 - 6x + 4$ ***which will produce the phantom graphs by substituting*** z^2 ***into*** $y = x^3 - 3xz^2 - 3(x^2 - z^2) + 4x - 2$
This is what the whole graph is like showing roots $x = 1, 1 + i$ and $1 - i$ where the graphs cross the complex x plane:



I found that when there are "missing" *y* values on a graph, there must be some *phantom graphs* which "hide" the complex solutions.

One of my favourites is the hyperbola $y^2 = x^2 + 25$



Notice that y values are ≥ 5 or ≤ -5 but there are **apparently** none between 5 and – 5.

Example if $y = 7$ we can find the two x values:

If $49 = x^2 + 25$

$24 = x^2$

$x \approx \pm 4.9$ see graph.

However, we can find x values for $-5 \leq x \leq 5$

If $y = 4$ then $16 = x^2 + 25$ $-9 = x^2$ $x = \pm 3i$	If $y = 3$ then $9 = x^2 + 25$ $-16 = x^2$ $x = \pm 4i$	If $y = 0$ then $0 = x^2 + 25$ $-25 = x^2$ $x = \pm 5i$

Of course, these are points on a CIRCLE of radius 5 units, in the plane perpendicular to the hyperbola!



Now there are no "missing" y values because the graph will cross any horizontal plane exactly two times.

By far the most challenging complex algebra was needed in finding the equations of the *phantoms* for the function $y = \frac{x^4}{x^2 - 1}$



We can see that there are no real values of y in the interval $0 \leq y \leq 4$

Interestingly, if we consider a general y value such as $y = c$ we get $\frac{x^4}{x^2 - 1} = c$ which produces a typical quartic equation $x^4 - cx^2 + c = 0$ which of course has 4 solutions.

However, if we draw $y = 5$ (see below) it crosses 4 times so the equation $\frac{x^4}{x^2 - 1} = 5$ clearly has 4 real solutions because it crosses the graph twice.



If we draw $y = 2$ there are no real solutions because the line does not intersect any part of the curve.

If we draw $y = -2$ it crosses TWO times so now there are apparently only **two** real solutions.

But when we consider the graph with its Phantoms, we see that **any horizontal plane $y = c$** crosses the graph 4 times. This further verifies the truth of the Fundamental Theorem of Algebra.



Incidentally, the equation of the bottom purple phantom is: $y = 1 - z^2 - \dfrac{1}{z^2 + 1}$

and the equation of the top blue phantom is: $y = x^2 - z^2 + 1 + \dfrac{(x^2 - z^2 - 1)}{(x^2 - z^2 - 1)^2 + 4x^2z^2}$

Another graph which has missing y values is the exponential graph.
This idea had never occurred to me before.
The basic 2D graph of $y = e^x$ is below:



Because e^x is always > 0 I never thought there were any y values **under** the x axis.
But then I thought, **"Why not?"**

I approached the problem again by allowing complex x values and investigated if it were possible to produce more real y values.

I let $y = e^{x+iz} = e^x . e^{iz} = e^x(\cos z + i \sin z)$ using Euler's formula.

If $z = 0$ we just get the basic curve $y = e^x$

If $z = 2n\pi$ (i.e. any multiple of 360^0) we get:
$y = e^x(\cos 2n\pi + i \sin 2n\pi)$
$= e^x (\quad 1 \quad + 0i \quad) = e^x$
This means there are infinitely many versions of $y = e^x$ all parallel to the basic 2D curve spaced at intervals of 2π in the imaginary x direction.

If $z = (2n+1)\pi$ (ie multiples of $180 + 360^0$) we get:
$y = e^x(\cos(2n+1)\pi + i \sin(2n+1)\pi)$
$= e^x (-1 \quad + \quad 0i \quad) = -e^x$
This means there are infinitely many versions of $y = e^x$ but **upside down** and all parallel to each other starting at $\pm\pi$ and then spaced at intervals of 2π to each other in the imaginary x direction.



I also thought that the basic trigonometrical graphs such as $y = sin(x)$ and $y = cos(x)$ only existed between 1 and –1.

It seemed nonsense that $sin(x)$ could equal 2 or any value > 1 or < -1.

This is the basic $y = sin(x)$ graph in 2D



I approached the problem again by allowing complex x values and investigated if it were possible to produce more real y values.

I let $y = sin(x + iz)$

$= sin(x)\ cos(iz) + cos(x)\ sin(iz)$

$= sin(x)\ cosh(z) + cos(x)\ sinh(-y)$

$= sin(x)\ cosh(z) - cos(x)\ sinh(z)$

If $z = 0$ then we just get the basic 2D curve $y = sin(x)$

If $x = \frac{\pi}{2} + 2n\pi$ then $y = sin(\frac{\pi}{2} + 2n\pi + iz) = 1 \times cosh(z) - 0 \times sinh(z) = cosh(z)$

This means at every maximum point a phantom $y = cosh(z)$ appears.

If $x = -\frac{\pi}{2} - 2n\pi$ then $y = -1 \times cosh(z) - 0 \times sinh(z) = -cosh(z)$

This means at every minimum point a phantom $y = -cosh(z)$ appears.



Here we can see that any equation ***sin(x) = c*** will have infinitely many solutions.
If we restrict our domain to just one period where $0 \leq x \leq 2\pi$ then equations like

sin(x) = ½ will just have 2 solutions and ***sin(x) = 2*** will also have 2 solutions.



There are some graphs which do not have *phantoms*, for example: $y = \frac{1}{x}$



Notice that any horizontal line will always cross this curve **one time.**
There are no "missing" values which is the usual sign that there are "*phantoms*".
This is because if we solve $\frac{1}{x} = c$ we get cx = 1 so $x = \frac{1}{c}$ for which there is always just one x value for all values of c except $c = 0$ which is the horizontal asymptote of course.

If we tried to find any phantoms by algebra we would do the following:
Let $y = \frac{1}{x + iz} = \frac{1}{x + iz} \times \frac{(x - iz)}{(x - iz)} = \frac{x}{x^2 + z^2} - \frac{iz}{x^2 + z^2}$

We would be looking for complex x values which produce only real y values so the imaginary part of y has to be zero.
This means $\frac{iz}{x^2 + z^2} = 0$ **so** $z = 0$ which simply means that $y = \frac{1}{x + 0i} = \frac{1}{x}$
which is just the original equation.

However considering $y = \frac{1}{x^2}$ it is obvious that there are "missing y values" below the x axis.



Here we see that any horizontal line $y = c$ where $c > 0$ will cross this graph twice. This is because the equation $\frac{1}{x^2} = c$ becomes $x^2 = c$ which always has two solutions.

To find the equation of the phantom graphs for $y = \frac{1}{x^2}$ we follow the same method as above:

Let $y = \frac{1}{(x + iz)^2} = \frac{1}{(x + iz)^2} \times \frac{(x - iz)^2}{(x - iz)^2}$

$$y = \frac{x^2 - z^2 - 2xzi}{(x^2 + z^2)(x^2 + z^2)}$$

$$\text{So } y = \frac{x^2 - z^2}{(x^2 + z^2)^2} - \frac{2xzi}{(x^2 + z^2)^2}$$

We would be looking for complex x values which produce only real y values so the imaginary part of y has to be zero.

This means $\frac{2xzi}{(x^2 + z^2)} = 0$

So that either $z = 0$ which just implies $y = \frac{1}{x^2}$ which is the original equation

or $x = 0$ which means $y = \frac{0 - z^2}{(0^2 + z^2)^2} = \frac{-1}{z^2}$ which produces the purple phantoms as shown on the graph below:



Now we can see that any horizontal plane $y = c$ except $c = 0$ will cross this graph twice.

NB
To draw a graph with ***real values of x and y*** we need **2 axes** and the graph is in **2 Dimensions**.

To draw a graph where ***y is real*** but ***x can have a real part and an imaginary part*** (like $3 + 4i$) we will need **3 axes** namely a ***real y axis, a real x axis and an imaginary x axis.*** This requires **3 Dimensions**.

To draw a graph where **x and y can both have a real part and an imaginary part** we would need **4 axes**. **This requires 4 Dimensions which I can't draw!**